



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : C12N 15/12, 5/10, C07K 14/475, 16/22, C12Q 1/68, G01N 33/574, A61K 31/70</p>	<p>A2</p>	<p>(11) International Publication Number: WO 97/12972 (43) International Publication Date: 10 April 1997 (10.04.97)</p>
<p>(21) International Application Number: PCT/IB96/01113 (22) International Filing Date: 30 September 1996 (30.09.96) (30) Priority Data: 9519928.7 29 September 1995 (29.09.95) GB 9612368.2 13 June 1996 (13.06.96) GB (71) Applicant (for all designated States except US): UNIVER- SITA' DEGLI STUDI DI SIENA [IT/IT]; Via Banchi di Sotto, 55, I-53100 Siena (IT). (72) Inventor; and (75) Inventor/Applicant (for US only): OLIVIERO, Salvatore [IT/IT]; Via del Giglio, 20, I-53100 Siena (IT). (74) Agent: HALLYBONE, Huw, George; Carpmaels & Ransford, 43 Bloomsbury Square, London WC1A 2RA (GB).</p>		<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p>Published Without international search report and to be republished upon receipt of that report.</p>
<p>(54) Title: REGULATED GENES AND USES THEREOF</p> <p>(57) Abstract</p> <p>A nucleotide molecule encoding a protein encoded by a Fos regulated gene or a fragment thereof, wherein said protein or fragment thereof is encoded by any one of the nucleotide sequences shown in Figure 1 or 2 or a fragment thereof, including allelic variants and species variants of the nucleotide sequences.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic			SE	Sweden
CG	Congo	KR	Republic of Korea	SG	Singapore
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LR	Liberia	SZ	Swaziland
CS	Czechoslovakia	LT	Lithuania	TD	Chad
CZ	Czech Republic	LU	Luxembourg	TG	Togo
DE	Germany	LV	Latvia	TJ	Tajikistan
DK	Denmark	MC	Monaco	TT	Trinidad and Tobago
EE	Estonia	MD	Republic of Moldova	UA	Ukraine
ES	Spain	MG	Madagascar	UG	Uganda
FI	Finland	ML	Mali	US	United States of America
FR	France	MN	Mongolia	UZ	Uzbekistan
GA	Gabon	MR	Mauritania	VN	Viet Nam

REGULATED GENES AND USES THEREOF

The present invention relates to the nucleotide sequences of Fos regulated genes, the proteins encoded by the sequences, 5 uses of the sequences and encoded proteins, and transgenic animals comprising one or more of the sequences. The present invention also relates to antibody molecules having affinity for the encoded proteins and uses of the antibody molecules, and antisense nucleotide molecules and uses of 10 the antisense nucleotide molecules.

The transcription factor AP-1 is involved in a number cellular processes, including cell proliferation, differentiation and neuronal function (see Angel and Karin, 15 1991).

AP-1 is considered to exert its effect by binding to a DNA recognition sequence, known as the AP-1 element, found in the promoter and enhancer regions of genes. The AP-1 20 element has the consensus sequence in TGA G/C TCA.

A number of genes have been found which contain AP-1 elements in their regulatory regions including c-Jun (Angel et al., 1988), MCP-1 (Rollins et al., 1988), Stromelysin 25 (Kerr et al., 1988), Type I collagenase (Schonthal et al., 1988) and Interleukin II (Farrar et al., 1989).

AP-1 is composed of dimeric complexes formed between Jun (c-Jun, Jun-B and Jun D) and Fos (c-Fos, Fos B, Fra-1 and Fra- 30 2) proteins. The Fos component of AP-1 has been found to be the limiting component of AP-1 activity in cycling cells (see Kovary and Bravo, 1991).

c-Fos is a nuclear proto-oncogene which has been implicated 35 in a number of important cellular events, including a cell proliferation (Holt et al., 1986; Riabowol et al., 1988), differentiation (Distel et al., 1987; Lord et al., 1993 and tumorigenesis (Curren et al., 1983; Miller et al., 1984;

Ruther et al., 1989).

c-Fos encodes a 62kDa protein which forms heterodimers with c-Jun, forming an AP-1 transcription factor which binds to DNA at an AP-1 element and stimulates transcription.

Fos gene products can also repress gene expression. Sassone et al. (1988) showed c-Fos inhibits its own promoter and Gius et al. (1990) and Hay et al. (1989) showed c-Fos inhibits early response genes Egr-1 and c-myc.

AP-1 factors have also been shown to inhibit expression of the MHC class I and PEPCK genes (see Gurney et al., 1992 and Howcroft et al., 1993).

15

It can therefore be seen that Fos regulated genes are extremely important for the correct expression of genes which lead to changes in the cell phenotype. The importance of Fos genes was clearly demonstrated by generating mice deficient in c-Fos (see Hu et al., 1994). The c-Fos deficient mice were viable, but displayed a range of tissue-specific developmental defects, including osteopetrosis, delayed gametogenesis and lymphopenia and behavioural abnormalities.

25

The c-Fos deficient mice were used to generate fibroblast cell lines and the expression of two genes was found to be abnormally low. The two genes were Stromelysin and Type I collagenase. Both genes were previously identified as having AP-1 sites in their regulatory sequences (see Kerr et al., 1988 and Schonthal et al., 1988).

Stromelysin and Type I collagenase have been implicated in embryonic tissue development (Brenner et al., 1989), injured tissue remodelling (Hasty et al., 1990; Woessner and Gunja, 1991) and in tumour progression and metastasis (Liotta and Stetler, 1990).

Superti-Furga et al., (1991) showed that c-Fos activity can be controlled hormonally by fusing the mouse c-Fos protein to the ligand binding domain of the human estrogen receptor. The fusion protein was found to stimulate AP-1 dependant transcription in a strictly hormone-dependant manner. Using the fusion protein an AP-1 regulated gene, Fit-1, was found. Fit-1 was found to encode a secreted or membrane bound protein depending on the splicing pattern.

- 10 The present invention relates to the nucleotide sequences encoding two novel Fos regulated genes.

The present invention provides a nucleotide molecule encoding a protein encoded by a Fos regulated gene or a fragment thereof, wherein said protein or fragment thereof is encoded by a nucleotide sequence shown in Figure 1 or 2, or a fragment thereof, including allelic variants and species variants of the nucleotide sequences.

- 20 The term "nucleotide molecule" used herein refers to nucleotides of any length, either ribonucleotides or deoxyribonucleotides. The term encompasses both double and single stranded molecules. It also includes known types of modifications, for example labels which are known in the art, methylation, "caps", substitution of one or more of the naturally occurring nucleotides with an analog, internucleotide modifications such as, for example those with uncharged linkages (e.g., methyl phosphonates, phosphotriesters, phosphoamidates, carbamates, etc.) and with charged linkages (e.g., phosphorothioates, phosphorodithioates, etc.), those containing pendant moieties, such as, proteins (including nucleases, toxins, antibodies, signal peptides, poly-L-lysine, etc.), those containing intercalators (e.g., acridine, psoralen, etc.), those containing chelators (e.g., metals, radioactive metals, boron, oxidative metals, etc.), those containing alkylators and those containing modified linkages (e.g., alpha anomeric nucleic acids, etc.).

The nucleotide molecule of the present invention may encode the protein of a Fos regulated gene or a fragment thereof.

5 The term "fragment" used in relation to the proteins refers to fragments which are of sufficient length to be unique to the presently claimed protein (e.g., 10, 15, 20 or 25 consecutive amino acids in length). Preferably, the protein fragments are capable of eliciting at least part of an
10 activity of the full protein. Particularly preferred fragments comprise a conserved region of a gene which has been found to be homologous with a number of other genes. Such conserved regions are considered to have a specific function.

15

The nucleotide sequences shown in Figures 1 and 2 will, as with most naturally occurring nucleotide sequences, have a number of other forms, such as allelic variants and species variants. Such variants and any other naturally occurring
20 forms of the nucleotide sequences of the present invention are also considered to form a part of the present invention. Such variants should have at least 60%, preferably 80% and most preferably 90% sequence homology with the sequences shown in figure 1 or 2 or fragments thereof.

25

The present invention also relates to the nucleotide molecule of the present invention wherein the protein or a fragment thereof encoded by the sequence shown in Figure 1 or 2 or a fragment thereof is altered.

30

Preferred altered proteins or fragments thereof, are those that still retain their activity and preferably have a homology of at least 80%, more preferably 90% and most preferably 95% to the protein or a fragment thereof encoded
35 by the sequence shown in Figure 1 or 2 or a fragment thereof. Preferably such altered proteins or fragments thereof differ by only 1 to 10 amino acids. It is further preferred that the amino acid changes are conservative.

- Conservative changes are those that replace one amino acid with one from the family of amino acids which are related in their side chains. For example, it is reasonable to expect that an isolated replacement of a leucine with an isoleucine or valine, an aspartate with a glutamate, a threonine with a serine, or a similar conservative replacement of an amino acid with a structurally related amino acid will not have a major effect on the biological activity of the protein.
- However, it is sometimes desirable to alter amino acids in order to alter the biological activity of the protein. For example, mutations which abolish or enhance one or more of the functions of the protein can be particularly useful. Such mutations can generally be made by altering any conserved sequences of protein. Mutations which increase the number of amino acids which are capable of forming disulphide bonds with other amino acids in the protein are particularly preferred in order to increase the stability of the protein. Mutations which decrease the number of amino acids which are capable of forming disulphide bonds with other amino acids in the protein may also be made if it is desired to decrease the stability of the protein. It is preferred that such altered proteins or fragments thereof have a homology of at least 80%, more preferably 90% and most preferably 95% to the protein or a fragment thereof encoded by the sequence shown in Figure 1 or 2 or a fragment thereof. Preferably such altered proteins or fragments thereof differ by only 1 to 10 amino acids.
- The nucleotide molecule of the present invention can be obtained by methods well known in the art. For example, the sequences may be obtained by genomic cloning or cDNA cloning from suitable cell lines or from DNA or cDNA derived directly from the tissues of an organism, such as a mouse. Suitable cell lines include any fibroblast cell lines such as the 3T3 cell line, described by Hu et al., (1994). Positive clones may be screened using appropriate probes for the nucleotide molecule desired. PCR cloning may also be

used. The probes and primers can be easily generated given that the sequences encoding the protein or a fragment thereof encoded by the nucleotide molecule of the present invention are given herein.

5

Numerous standard techniques known in the field of molecular biology may be used to prepare the desired nucleotide molecules or the probes and primers for identifying the positive clones. The nucleotide molecules probes or primers
10 may be synthesised completely using standard oligonucleotide synthesis methods, such as the phosphoramidite method.

Numerous techniques may be used to alter the DNA sequence obtained by the synthesis or cloning procedures, and such
15 techniques are well known to those skilled in the art. For example, site directed metagenesis, oligonucleotide directed mutagenesis and PCR techniques may be used to alter the DNA sequence. Such techniques are well known to those skilled in the art and are described in the vast body of literature
20 known to those skilled in the art, for example Sambrook et al., (1989).

The present invention further provides the protein encoded by the nucleotide molecule of the present invention.

25

Preferably, the protein encoded by the nucleotide molecule of the present invention has the amino acid sequence shown in Figure 1 or 2, or a fragment thereof.

30 The term "protein" as used herein refers to a polymer of amino acids and does not refer to a specific length of the product; thus, peptides, oligopeptides and proteins are included within the term protein. The term also does not refer to or exclude post-expression modifications of the
35 protein, for example, glycosylations, acetylations and phosphorylations. Included in the definition are proteins containing one or more analogs of an amino acid (including for example, unnatural amino acids), proteins with

substituted linkages, as well as other modifications known in the art, both naturally occurring and synthesised.

The protein of the present invention can be obtained from
5 cells that naturally produce the protein such as fibroblast cells using standard purification techniques. However, it is preferred that a suitable host cell and vector system is used for the expression of the nucleotide molecule of the present invention. The nucleotide molecule of the present
10 invention can be expressed in a variety of different expression systems, for example, those used with mammalian cells, baculoviruses, bacteria and eukaryotic microorganisms such as yeasts.

15 All the above mentioned expression systems are known in the art and expressing nucleotide sequences is now a standard technique known all skilled in the art.

Preferably, eukaryotic, e.g. mammalian, host cell expression
20 systems are used. In particular, suitable mammalian host cells include chinese hamster ovary (CHO) cells, HeLa cells, baby hamster kidney (BKH) cells, cells of hepatic origin such as HepG2 cells, and myeloma or hybridoma cell lines.

25 The present invention further provides a vector for the expression of the nucleotide molecule of the present invention, comprising a promoter and the nucleotide molecule of the present invention.

30 A mammalian promoter can be any DNA sequence capable of binding mammalian RNA polymerase and initiating the downstream transcription of a coding sequence into mRNA. Particularly useful promoters are those derived from mammalian viral genes, such as the SV40 early promoter,
35 adenovirus major late promoter and the herpes simplex virus promoter. Additionally, sequences from non-viral genes can also be used as promoters, such as from the murine metallothionein gene.

The nucleotide molecule of the present invention may be expressed intracellularly in mammalian cells. A promoter sequence may be directly linked with the nucleotide molecule
5 of the present invention, in which case the first amino acid at the N-terminus of the encoded protein will be a methionine encoded by the start ATG codon.

Alternatively, the protein encoded by the nucleotide
10 molecule of the present invention can be secreted from the cell by linking a nucleotide sequence encoding a leader sequence to the nucleotide molecule of the present invention. The encoded fusion protein will comprise a leader sequence fragment and the protein encoded by the
15 nucleotide molecule of the present invention. The leader sequence will lead to the secretion of the fusion protein out of the cell. Preferably, there are processing sites between the leader sequence and the protein encoded by the nucleotide molecule of the present invention allowing the
20 leader sequence to be cleaved off enzymatically or chemically. An example of such a leader sequence is the adenovirus tripartite leader.

The vector of the present invention is preferably a nucleic
25 acid vector comprising DNA. The vector may be of linear or circular configuration and can be adapted for episomal or integrated existence in the host cell, as set out in the extensive body of literature known to those skilled in the art. The vectors may be delivered to cells using viral or
30 non-viral delivery systems. The choice of delivery system will determine whether the DNA molecule is to be incorporated into the cell genome or remain episomal.

The vector of the present invention can comprise further
35 control elements such as polyadenylation signals, transcription termination signals, enhancers, locus control regions (LCRs), etc.

The present invention further provides a host cell transformed with the vector of the present invention.

Transformation refers to the insertion of an exogenous polynucleotide into a host cell, irrespective of the method used for the insertion, for example, direct uptake, transduction, f-mating or electroporation. The exogenous polynucleotide may be maintained as a non-integrated vector (episome), or may be integrated into the host genome.

10

Preferably, the host cell is a eukaryotic cell, more preferably a mammalian cell, such as chinese hamster ovary (CHO) cells, HeLa cells, baby hamster kidney (BKH) cells, cells of hepatic origin such as HepG2 cells, and myeloma or hybridoma cell lines.

The present invention further provides a method for producing the protein encoded by the nucleotide molecule of the present invention, comprising transfecting a host cell with the vector of the present invention, culturing the transfected host cell under suitable conditions in order to lead to the expression of the DNA molecule and the production of the desired protein. The protein may then be harvested from the transfected cells or from the cell growth media, depending on whether the protein is secreted, using standard techniques.

20
30

The present invention further provides the nucleotide molecule of the present invention for use in therapy.

The present invention further provides the use of the nucleotide molecule of the present invention in the manufacture of a composition for the treatment of developmental disorders.

35

The present invention further provides the use of the nucleotide molecule of the present invention in the treatment of developmental disorders.

Fos regulated genes are known to be involved in development and cell differentiation. Accordingly, the discovery of genes regulated by Fos has implications in the control of development and cell differentiation.

The nucleotide sequences shown in Figure 1 and Figure 2 have been found to have a similar sequence to genes of a family of growth factors characterised by the Platelet Growth Factor (PDGF) family signature. The most clearly related sequence is that of the Vascular Endothelial Growth Factor (VEGF). VEGF forms a homodimer which is a growth factor active in angiogenesis and endothelial cell growth (see Keck et al., 1989 and Leung et al., 1989). VEGF has also been used to stimulate angiogenesis and thereby produce a therapeutic effect (see Takeshita et al., 1994).

The protein encoded by the sequence in Figure 1 is a mouse protein and the protein encoded by the protein in Figure 2 is the human homologue of the mouse protein encoded by the sequence given in Figure 1. Both the proteins are herein referred to as c-Fos Induced Growth Factor (FIGF).

The use of the nucleotide molecule of the present invention in therapy can therefore be seen to be an important application of the sequences of the Fos regulated genes of the present invention.

The nucleotide sequences shown in Figure 1 and Figure 2 are of particular interest in lung disorders as it has been found that the nucleotide sequences are mainly expressed in the lungs. Particular lung disorders which may be treatable using the nucleotide molecule encoding the protein or fragments thereof which are encoded by the sequence shown in Figure 1 or Figure 2, include pneumonia and pneumoconiosis. The nucleotide molecule may also be used following pneumonectomy in order to aid in lung re-growth.

The nucleotide sequence in Figure 2 has been mapped to human chromosome Xp22, near the locus that maps for the pathology spondyloepiphyseal dysplasia (SEDL). The genetic map of this region is described by Ferrero et al. (1995) and the mapping of the SEDL disease is described by Heuertz et al. (1993). SEDL may therefore be treatable using the nucleotide molecule encoding the protein or fragments thereof, which are encoded by the sequence given in Figure 1 or in Figure 2.

10

As previously discussed, Fos regulated genes have been found to be involved in tumour progression and metastasis. By inhibiting Fos regulated genes it is possible to inhibit or suppress tumour growth.

15

Previously Kim et al., (1983) suppressed tumour growth by injecting monoclonal antibodies specific for VEGF. As stated previously, VEGF has a similar nucleotide sequence to the nucleotide sequences shown in Figure 1 and Figure 2.

20 Accordingly, by inhibiting either the *in vivo* expression, translation, etc. of the native nucleotide molecules of the present invention, tumour growth may be inhibited or suppressed.

25 The actions of the Fos regulated genes corresponding to the nucleotide molecules of the present invention may be inhibited by a number of techniques. Preferred techniques include antisense based techniques, ribozyme based techniques, and antibody based techniques.

30

Antibody molecules having specificity for the protein encoded by the nucleotide molecules of the present invention can be used to block the function of the protein and thereby inhibit or suppress tumour growth.

35

The present invention further provides antibody molecules having specificity for the protein of the present invention.

The antibody molecules may be a complete polyclonal or monoclonal antibody or antigen binding fragments, such as Fv, Fab, F(ab')₂ fragments and single chain Fv fragments thereof. The antibody molecule may be a recombinant
5 antibody molecule such as a chimeric antibody molecule preferably having human constant regions and mouse variable regions, a humanised CDR grafted antibody molecule or fragments thereof. Methods for producing such antibodies are well known to those skilled in the art and are described
10 in EP-A-0120694 and EP-A-0125023.

The present invention further provides the antibody molecule of the present invention for use in therapy.

15 The present invention also provides the use of the antibody molecule of the present invention in the manufacture of a composition for the treatment of proliferative diseases such as cancer.

20 The present invention further provides the use of the antibody molecule of the present invention for the treatment of proliferative diseases such as cancer.

The present invention further provides an antisense
25 nucleotide molecule or a fragment thereof, having the complementary sequence to the nucleotide molecule or a fragment thereof, of the present invention.

The antisense nucleotide molecule of the present invention
30 can be generated using the same standard techniques as for the nucleotide molecule of the present invention.

The present invention further provides an antisense vector for the expression of the antisense nucleotide molecule of
35 the present invention comprising a promoter and the antisense nucleotide molecule.

The antisense vector is identical to the nucleic acid vector

of the present invention except that the vector contains the antisense nucleotide molecule of the present invention.

The present invention further provides the antisense vector
5 of the present invention for use in therapy.

The present invention further provides the use of the antisense vector of the present invention in the manufacture of a composition for the treatment of cell proliferative
10 diseases such as cancer.

The present invention further provides the use of the antisense vector of the present invention in the treatment of cell proliferative diseases such as cancer.
15

The present invention further provides a vector for the expression of a ribozyme, comprising a promoter and a nucleotide sequence encoding a ribozyme capable of cleaving the RNA transcript of the nucleotide molecule of the present
20 invention.

The vector encoding the ribozyme is identical to the vectors previously described except that the vector encodes a ribozyme. The ribozyme being capable of cleaving the RNA
25 transcript of the nucleotide molecule of the present invention. Techniques for producing such ribozymes are known to those skilled in the art and are discussed in Cantor et al., (1993).

30 The present invention further provides the ribozyme encoding vector of the present invention for use in therapy.

The present invention further provides the use of the ribozyme encoding vector of the present invention in the
35 manufacture of a composition for the treatment of cell proliferative diseases such as cancer.

The present invention further provides the use of the

ribozyme encoding vector of the present invention in the treatment of cell proliferative diseases such as cancer.

5 A further object to the present invention is the use of the protein of the present invention in identifying the receptor or receptors of the protein or of a protein complex comprising the protein.

10 Methods for identifying receptors are well known to those skilled in the art and have been widely described in the literature. However, basically there are three major ways of identifying receptors:

- 15 i. Test all known receptors that bind to similar molecules. This is particularly useful for the protein encoded by the DNA sequences shown in Figure 1 and Figure 2, as VEGF has been found to have a similar sequence.
- 20 ii. Perform a binding purification step. For example, the protein of the present invention or a protein complex comprising the protein of the present invention can be immobilised on to a solid support and numerous possible receptor molecules, especially membrane proteins, are then passed over the solid support. A binding
25 purification procedures is described in Schusted et al., (1995).
- 30 iii. By screening expression libraries in order to find cells lacking the receptor or receptors and then utilising the receptor cloning method described by Seed and Aruffo, (1987).

Other methods are also known to those skilled in the art and can be used in order to find the receptor or receptors.
35

On identifying the receptor or receptors it will be possible to design drugs that block or enhance the activity of the receptor or receptors and produce antibody molecules that

block the receptor or receptors. Once the DNA sequence of the receptor or receptors are known, a number of gene therapies may be designed for correcting errors in the receptor or receptors, or for blocking expression of the
5 receptor or receptors.

The present invention further provides the use of the protein of the present invention in an assay for identifying antagonists or agonists of the protein which may be used as
10 drugs in the treatment of cancer and developmental disorders respectively. Assays for identifying such potential drugs are frequently used and are well known to those skilled in the art. An example of such an assay is clearly described in Tsunoda et al., (1994).

15 The present invention further provides the use of the nucleotide molecule, antisense nucleotide molecule, protein or antibody molecule of the present invention or any combination thereof, in diagnosing a pathological state or
20 a predisposition to a disease.

The nucleotide molecule or antisense nucleotide molecule of the present invention may be used in determining the presence of the gene corresponding to the nucleotide
25 molecule or in determining the amount of RNA transcribed from the gene.

The protein of the present invention may be used in an assay for determining the amount of protein encoded by the gene
30 corresponding to the nucleotide molecule of the present invention.

The antibody molecule of the present invention may be used in an assay for determining the amount of protein encoded by
35 the gene corresponding to the nucleotide molecule of the present invention. An example of an assay for determining the amount of protein using the antibody molecule of the present invention is a competitive binding assay.

By determining the presence of the gene corresponding to the nucleotide molecule of the present invention or the transcribed RNA or the protein encoded by the gene, it is possible to diagnose a pathological state or a predisposition to a disease caused by the presence of the gene of the over expression of the gene.

The present invention further provides the use of the nucleotide molecule of a present invention in the generation of transgenic animal. In particular, the invention provides the use of such nucleotide molecules for the generation of non-human transgenic animals, especially transgenic mice.

Transgenic animals can be generated which are suitable as models for research. For example, transgenic animals which overexpress the nucleotide molecule of the present invention could be used in order to determine what effects overexpression will have. Alternatively, transgenic animals can be generated where the native nucleotide molecule of the present invention is "knocked out". The effect of "knocking out" the nucleotide molecule could then be investigated.

Methods for generating such transgenic animals are well known to those skilled in the art and can be easily performed given that the nucleotide molecules to be over expressed or "knocked out" are disclosed herein.

The transgenic animals of the present invention could also be subsequently bred with either Fos over expression mice or Fos "knocked out" mice, in order to determine the effects of altered Fos control.

The present invention also provides a nucleotide molecule comprising all or part of the sequence shown in any one of Figures 1 or 2.

The nucleotide molecule comprising all or part of the

sequence shown in any one of Figures 1 or 2 may encode a protein or may be non-coding. Preferably, the nucleotide molecule additionally encodes the control sequences of the Fos gene corresponding to the nucleotide sequence shown in
5 any one of Figures 1 or 2. It is further preferred that the nucleotide molecule encodes a sequence which confers Fos regulation to a gene. It is particularly preferred that the nucleotide molecule comprises the sequence TGACTCA.

10 The present invention is now illustrated in the appended examples with reference to the following figures.

Figure 1

DNA sequence of Fos regulated gene F0401, showing the
15 encoded protein sequence and the regions homologous to VEGF (underlined).

Figure 2

DNA sequence of Fos regulated gene HF175 (human homologue of
20 F0401), showing the encoded protein.

Figure 3

Alignment of the protein encoded by FIGF with the conserved domain of the PDGF/VEGF family of growth factors. Dots
25 indicate the cysteine residues which are characteristic of these growth factors.

Figure 4

(A) Immunoprecipitation assay of the FIGF protein. COS-7
30 cells transfected with the vector alone (-) or with a vector containing the FIGF coding sequence under the control of a CMV promoter (+) were metabolically labelled for 1 hour with [³⁵S]Methionine and [³⁵S]Cysteine each at a concentration of 100 µCi/ml. After 1 hour or 22 hours chase, conditioned
35 media and cell lysates were immunoprecipitated separately with anti-FIGF polyclonal antibodies. [The FIGF protein was expressed in E. coli under the control of the T5 promoter. The cDNA fragment, from the coding region of FIGF,

was generated by PCR from the Methionine residue at position +40 and cloned into the pQE-31 vector (Qiagen) to obtain a fusion protein with a N-terminal Histidine tag. The protein was expressed in TG1 bacteria (pREP+) by induction for 4 hours at 37 °C in the presence of 2mM isopropyl-b-D-thiogalactopyranoside. The recombinant protein was exclusively localized in inclusion bodies, and was purified on a column of Ni-NTA-Resin, under denaturing conditions according to the manufacturer's protocols (Qiagen). Antibodies were raised by injecting rabbits with 200 µg of recombinant FIGF in form of denaturated protein in complete Freund's adjuvant. Serum was prepared after 4 injections in incomplete Freund's adjuvant at 3-week intervals]. The immunocomplexes were collected by protein-A Sepharose beads (Pharmacia) and separated on 12% SDS-PAGE in the presence of 3% b-mercaptoethanol. Arrows indicate specific bands present only in FIGF transfected cells.

(B) Mitogenic activity measured as [3H]-thymidine incorporation in c-fos (-/-) fibroblasts. Cells were incubated with conditioned medium of COS-7 cells transfected with the FIGF expression vector or with the vector alone. One day after transfection the cells were split and kept in 2% serum. Conditioned media were collected after 120 hours.

(C) Mitogenic activity measured as [3H]-thymidine incorporation in c-fos (-/-) fibroblasts. Cells were incubated with conditioned media obtained from c-fos (-/-) stable clones, named FH-10.2, FH-10.5, FH-9.3, FH-9.6, FH-10.9 and c-fos (-/-) cells (mock), constitutively expressing exogenous FIGF under the control of the CMV promoter. Conditioned media were collected from cells cultured for 48 hours in 0.5% serum.

(D) Mitogenic activity measured as [3H]-thymidine incorporation in c-fos (-/-) fibroblasts. Cells were incubated with partially renatured recombinant FIGF. Under the same conditions, incubation with PDGF-BB (Sigma), used as a positive control, induces about 30 % higher thymidine incorporation, while VEGF (Sigma) does not induce incorporation above the background. The data shown are the

mean of six experiments performed with two different FIGF preparations.

(E) Mitogenic activity measured as [3H]-thymidine incorporation on mouse embryo fibroblasts. Cells were
5 incubated with partially renatured recombinant FIGF. MEF cells were obtained from 13-15 day embryos of B6D2F1 mice. The embryos were sacrificed, rinsed and trypsinized for 30 min at 37°C. The MEF cells were grown 24 hours in medium containing 0.5 % serum before addition of the growth
10 factors. Under the same conditions, incubation with PDGF-BB (Sigma), used as a positive control, induces about 30 % higher thymidine incorporation, while VEGF (Sigma) does not induce incorporation above the background. The data shown are the mean of six experiments performed with two different
15 FIGF preparations. The background values were subtracted in each experiment.

Figure 5

(A) Expression of FIGF in cultured cells. Northern blot
20 analysis of total RNA obtained from: c-fos (-/-) fibroblasts (lanes 1 - 3); a stable cell line, obtained from c-fos (-/-) cells, expressing exogenous c-fos (lanes 4 - 6), c-fos (+/+) fibroblasts (lanes 7 - 9). Cellular RNA was extracted by the guanidine thiocyanate method after incubation of cells
25 for 48 hours in 0.5 % serum (time 0). The serum concentration was increased to 10 % and total RNA was collected at 2 or 4 hours as indicated. Lanes 10 and 11 show FIGF expression in c-fos (-/-) fibroblasts transiently transfected with the vector alone (mock) or containing the
30 c-fos under the FBJ-LTR constitutive promoter (c-fos). The RNAs of the transiently transfected cells were collected 30 hours after culturing the cells in medium containing 0.5% serum. Each lane was loaded with 10µg of total cellular RNA.

35 (B) Expression of PDGF or VEGF. Total cellular RNAs from c-fos (-/-) cells (lanes 1-3) or from a stable cell line, obtained from c-fos (-/-) cells, expressing exogenous c-fos (lanes 4 - 6) were extracted as indicated in panel A.

Glyceraldehyde-phosphate-dehydrogenase (GAPDH) was used as a control for RNA loading.

Figure 6

- 5 Northern analysis of RNA poly A+ extracted from different mouse tissues.

Figure 7

- (A) Morphology of c-fos deficient cells. The cells were
10 stably transfected with the vector alone.
- (B) Morphology of a cell clone derived from c-fos deficient cells, stably transfected with the expression vector containing FIGF under the control of the CMV promoter.
- (C) Morphology of cells stably transfected with an
15 expression vector containing the FIGF cDNA in the antisense orientation under the control of the CMV promoter.
- (D) Morphology of cells stably transfected with the expression vector containing c-fos under the control of the FBJ-LTR promoter.
- 20 (E) A cell clone derived from the same cells as in D (expressing c-fos constitutively) transfected with an expression vector containing FIGF under the control of the CMV promoter.
- (F) A cell clone derived from the same cells as in D
25 (expressing c-fos constitutively) transfected with an expression vector containing the FIGF cDNA in the antisense orientation under the control of the CMV promoter.
- (G) c-fos (-/-) fibroblasts cultured for 120 hours in medium containing 0.5 % serum.
- 30 (H) Cells as in G but treated for 120 hours with partially renatured recombinant FIGF. Ten independent clones obtained from 3 independent transfections were analyzed. All showed morphological changes similar to those observed in the figure.

35

EXAMPLES

Cell Culture and Clone Isolations

Mouse fibroblast wild-type cells with respect to c-Fos

expression (+/+) and c-Fos-deficient (-/-) 3T3 cell lines and stably transfected cell line that constitutively express exogenous c-Fos were generated as described (Hu, et al., 1994). All cell lines were grown at 37°C with 5% CO₂ in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% fetal calf serum (FCS), glutamine and penicillin-streptomycin. Cells were cultured until reaching about 70% confluence, serum starved for 48 hours in DMEM containing 0.5% FCS and stimulated with DMEM containing 10% FCS for 0, 2 and 4 hours prior to RNA isolation. Total RNA was isolated using the guanidine-isothiocyanate method. mRNA differential display was performed as described by Laing et al. and modified by Bauer et al. (Bauer et al., 1993). Briefly, from the extracted RNA chromosomal DNA contamination was removed from 50 µg of the total RNA isolated by DNase I treatment. 0.2 µg of RNA, extracted at 2 or 4 hours after serum induction, was used for reverse transcription in a 40 µl reaction volume using dT₁₂mN primers and 300 U MMLV reverse transcriptase (Promega Corp., Madison, WI) with an incubation time of 60 min at 37°C. The PCR mixture for the cDNA amplification contained dT₁₂mN primer, one of the 20 10mer deoxyoligonucleotide primers with arbitrary sequence (Kit A - Operon Biotechnology Inc., Alameda, CA), ³³P-dATP (Amersham International plc, Buckinghamshire, England) and 1U Taq polymerase (Promega Corp.). Samples were subjected to 40 cycles of amplification using a PTC-100 thermocycler (MJ Research Inc., Watertown, MA). The cycling parameters were as follows: 94°C for 30 seconds, 42°C for 90 seconds, 72°C for 30 seconds and an additional extension period at 72°C for 10 min. 2 µl of the PCR mixture was adjusted with glycerol to 5% and loaded onto a 6% polyacrylamide gel without urea (Bauer et al., 1993). The bands of cDNA differentially expressed, were recovered from the gel and reamplified. Reamplified cDNA probes were run on a 1.5% agarose gel, purified and cloned into the pGEM-T vector using the TA cloning system (Promega Corp.) Positive clones were selected using the blue-white phenotype.

Characterisation and Sequencing of Novel Clones

Typically from one band we could obtain 1 to 3 different clones, which we utilized for the successive characterisation by Northern blot analysis. The cDNA fragments were labelled with ^{32}P -dCTP using a random primer labelling kit (Amersham International plc). Hybridisation signals were screened and quantitated by PhosphorImager using Image Quant software (Molecular Dynamics, Sunnyvale, CA). Plasmid DNA sequencing of cloned cDNA probes with either T7 or SP6 primer was carried out manually using the Sequenase V 2.0 Kit (US Biochemical Inc., Cleveland, Ohio). Briefly, the RNA extracted from the cells were subjected to amplification utilising random primers and the bands of a cell type are identified by comparison and isolated. The fragments obtained were tested in Northern blot with RNA from the cell lines to confirm that the corresponding mRNA are up regulated in Fos expressing cells. Then we generated our own cDNA library in lambda ZAP vectors from mouse fibroblasts cell lines to obtain the full length clones utilising a cDNA Synthesis and Cloning Kit (Stratagene). The screening was performed according to the manufacturer. Positive clones were first analysed by restriction map and the longest ones were subjected to DNA sequence.

Clone Analysis

F0401 sequence is shown in Figure 1 and the HF175 sequence is shown in Figure 2: a simple search analysis against the NIH and EMBL data banks revealed that F0401 and the human homologue FIGF are novel genes and their sequences are similar to the genes of a family of growth factors characterised by the Platelet growth Factor (PDGF) family signature. The consensus pattern of the family is: C-V-x(3)-R-C-x-G-C-C-N.

Members of this family form dimers with disulphide links and are potent mitogens. The most similar sequence to F0401 and HF175 is the Vascular Endothelial growth factor (VEGF) which forms an homodimer and is a growth factor active in

angiogenesis and endothelial cell growth (Keck et al., 1989; Leung et al., 1989). As VEGF is a growth factor its over expression can result in tumour angiogenesis (Plate et al., 1993). Recent reports indicate possible therapeutic use
5 based on VEGF inhibition in tumours (Kim et al., 1993) and on VEGF treatment to stimulate angiogenesis (Takeshita et al., 1994).

The following experiments were performed using F0401.

10

The FIGF predicted protein sequence has a hydrophobic sequence at the N-terminus which could code for a signal peptide. This long N-terminus region does not show significant homology to known proteins. However, there is
15 a positively charged domain in this region which may allow binding of the protein to the cell membrane or to the extracellular matrix.

To verify if FIGF is a secreted protein, we transfected
20 COS-7 cells with an expression vector containing the FIGF cDNA under the control of the cytomegalovirus (CMV) immediate early gene promoter. Polyclonal antibodies, raised against recombinant FIGF (as described previously), immunoprecipitated a specific band that is observed in both
25 the cell lysates and the conditioned media of the FIGF transfected COS-7 cells (Fig.4A). After 1 hour labelling followed by 1 hour chase a specific band was mainly present in the cell lysate while, after a chase longer than four hours, the protein accumulated in the cell supernatant.
30 Under non-denaturing conditions FIGF aggregated into a multimeric form. Addition of b-mercaptoethanol resulted in partial denaturation of the protein which migrated mostly as a 66 kDa band and only a minor fraction of the protein can be found as a monomer of the expected 33 kDa of
35 molecular mass (Fig.4A). These results show that FIGF is a secreted protein and can form dimers. Dimerization of FIGF could be predicted since the FIGF central domain is highly conserved and contains the cysteine residues involved in the

dimerization of both PDGF and VEGF. It was further investigated whether the conditioned medium of FIGF producing cells could promote cell growth in vitro, assayed as [3H]-thymidine incorporation (Vaziri et al. (1995)).

5 Conditioned medium was obtained either from transiently transfected COS-7 cells or from stable clones, derived from c-fos(-/-) fibroblasts, expressing FIGF under the control of the CMV promoter. The mitogenic activity of the medium containing FIGF was tested on c-fos(-/-) fibroblasts.

10 Conditioned medium from both transfected COS-7 (Fig. 4B) or stable fibroblast clones overexpressing FIGF (Fig. 4C) induces DNA synthesis in c-fos (-/-) fibroblasts. As in mammalian cells FIGF expression could induce the activation of other growth factors, which in turn would be responsible

15 for the [3H]-thymidine incorporation measured, we tested the mitogenic activity of a recombinant FIGF protein expressed in E. coli (as described previously). In order to obtain a biologically active recombinant protein, the purified FIGF protein from E. coli was partially renatured

20 in the presence of a mixture of reduced and oxidized glutathione. The purified recombinant protein was adjusted to 0.4 mg/ml and completely reduced in the presence of 8M Urea, 2% b-mercaptoethanol for 1 hour at 370 °C. The reduced protein was dialyzed against a solution containing 50 mM

25 Tris-Cl pH 8.0, 1M Urea, 5 mM reduced glutathione and 0.5 mM oxidized glutathione for 2 days, and against a solution containing 20 mM Tris-Cl pH 7.5, 0.7 M NaCl for 1 day, as described by Hoppe et al., Biochemistry, 28, pp 2956-2960 (1989); Hoppe et al., Eur. J. Biochem., 187, pp 207-214

30 (1990). The partially refolded recombinant FIGF induced DNA synthesis on c-fos (-/-) fibroblasts in a dose-dependent manner (Fig. 4D). As expected, c-fos(-/-) cells are also responsive to PDGF-BB, while the treatment with VEGF did not induce [3H]-thymidine incorporation in these cells. The

35 highest activity of DNA synthesis was obtained with 2 µg of purified FIGF. The apparently low specific activity of the recombinant FIGF observed, is most probably due to the low efficiency of FIGF correct refolding since FIGF contains 29

cysteine residues out of 358 amino acids. We also tested the mitogenic activity of the recombinant FIGF on mouse embryo fibroblasts (MEF). FIGF induced DNA synthesis on mouse embryo fibroblasts in a dose-dependent manner (Fig. 4E). The FIGF cDNA was isolated by differential screening of RNA from cells differing only for the expression of c-fos. Analysis of FIGF gene expression by Northern blot reveals that the FIGF messenger is barely detectable in c-fos (-/-) fibroblasts, while its expression is high in wild type c-fos (+/+) fibroblasts (Fig. 5A, compare lanes 1-3 with lanes 7-9). FIGF expression is completely restored in stable clones, derived from c-fos (-/-) cells, expressing exogenous c-fos under the control of the FBJ-LTR constitutive promoter (Hu et al. (1994)) (Fig. 5A, compare lanes 1-3 with lanes 4-6). The transient transfection of exogenous c-fos results in FIGF induction in c-fos (-/-) cells, although, due to the lower number of transfected cells, the induction observed is less pronounced (Fig. 5A lanes 10 and 11). Thus, FIGF expression is dependent on c-fos. Moreover, FIGF is not induced by the constitutive AP-1 yeast homologue GCN4. In mammalian cells GCN4 is able to activate most AP-1 target genes, but it is non-oncogenic. In wild type fibroblasts c-Fos is the major Fos protein associated with c-Jun or Jun B within the first hour after serum induction. Afterwards c-Fos is not detectable any longer and it is substituted by FraJ1 and FraJ2 in the AP-1 complex. In c-fos expressing cells, FIGF is highly expressed when cells are kept in low serum conditions and decreases to undetectable levels within six hours after serum induction (Fig. 5A). This pattern of FIGF expression can be observed both in wild type cells and in cells constitutively expressing c-fos (Fig. 5A). Thus, we observe a discrepancy between the expected peak of c-fos expression and the appearance of FIGF, whose messenger accumulates in the quiescent phase. The FIGF pattern of regulation suggests that, besides the expression of c-fos, additional regulatory controls are required for its activation. Although FIGF belongs to the PDGF/VEGF family of growth factors, its

expression is most similar to the expression of the growth arrest specific (gas) genes. Interestingly, one of them, gas6, acts as a growth factor. Both PDGF and VEGF growth factors are involved in tumour formation (Kim et al. 5 (1993)). Moreover, PDGF is the main serum mitogen which induces the transcription activation of c-fos. In order to compare the pattern of expression of these growth factor with respect to FIGF, we measured the PDGF and VEGF messengers levels in fibroblasts differing for the 10 expression of c-fos. As can be observed in Fig. 5B, the regulation of both PDGF and VEGF messengers is distinct from that of FIGF. These growth factors are rapidly induced following serum induction and their expression is independent of c-fos. Tumour progression is characterized 15 by morphological changes of the tumour that leads the mutated cells to loose their adhesion to the original neighbours and escape from the tissue of origin. c-fos has been implicated in tumour progression and its over-expression induces a transformed cell morphology in 20 fibroblasts and epithelial cells. As FIGF is a c-fos-dependent growth factor, it was analyzed whether its over-expression could induce fibroblast morphological transformation. As can be observed in Fig. 7, the constitutive expression of FIGF in fibroblasts induces a 25 transformed phenotype. Stable clones derived from c-fos (-/-) cells, constitutively expressing FIGF, acquire a spindle-shaped morphology, become more refractive, and detach from the plate (Fig 7, B versus A). On the contrary, stable clones expressing the FIGF antisense messenger 30 acquire a flat and less refringent phenotype (Fig. 7C), which is most similar to the phenotype of c-fos (-/-) cells kept in low serum conditions (Fig.7G). The over-expression of c-fos alters c-fos (-/-) cell morphology similarly to that observed with the over-expression of FIGF, although 35 the phenotype is less pronounced (Fig. 7D). The over-expression of both c-fos and FIGF leads to an extreme phenotype in fibroblasts: cells become longer, disorganized and lose contacts (Fig.7E). The expression of the FIGF

antisense messenger in cells constitutively expressing c-fos induces a reversion of the transformed phenotype (Fig. 7F). Thus, cells expressing c-fos but depleted of FIGF lose most of the transformed phenotype, suggesting that the morphology
5 observed in cells constitutively expressing c-fos is due to FIGF. Similar morphological alterations are also obtained by cell treatment with purified recombinant FIGF. c-fos (-/-) fibroblasts, kept in medium containing 0.5% serum for 120 hours stop growing, become flat large and less refractile
10 (Fig. 7G). Cell treatment with recombinant FIGF induces the refractile, elongated and non-adherent phenotype (Fig. 7H).

Tumours obtained from cells defective for c-fos cannot
15 undergo malignant progression even if they are carrying the activated v-H-Ras. Thus, the expression of c-fos is essential for the activation of target genes responsible for the malignant phenotype. FIGF is a c-fos-dependent autocrine growth factor able to induce cell division entry
20 and, when it is over-expressed, a transformed phenotype in fibroblasts. The data suggest that the role of c-fos in the activation of the malignant phenotype is due to the activation of FIGF.

25 Further experiments on FIGF using a probe specific for FIGF in Northern analysis of RNA derived from mouse tissues, show that the FIGF gene is only expressed in cells expressing Fos and poorly in cells that lack the Fos oncogene (Figure 5). The RNA blot used in the Northern assay was obtained from
30 Clontec. The analysis of its expression in the mouse tissues shows that FIGF is mainly expressed in lung (Figure 6) and is already present at day 7 of the mouse embryonal life (not shown).

35 FIGF is therefore a molecule related to the growth factor VEGF, positively regulated by the oncogene Fos. It could be implicated in tumours and in development.

References

1. Angel et al., (1988) Cell 55, pp 875-885
2. Angel and Karin, (1991), Biochim, Biophys. Acta, 1072,
5 pp 129-57
3. Bauer et al., (1993), NAR, 21, pp 4272-4280
4. Bergens et al., (1994), EMBO J., 13, pp 1176-1188
5. Brenner et al., (1989), Nature, 337, pp 661-663
6. Cantor et al., (1993), Proc. Natl. Acad. Sci. USA, 90,
10 pp 10932-10936
7. Curren et al., (1983), Mol. Cell. Biol., 3, pp 914-921
8. Distel et al., (1987) Cell, 49, pp 835-844
9. Farrar et al., (1989), Crit. Rev. Ther. Drug Carrier
Syst., 5, pp 229-261
- 15 10. Ferrero et al., (1995), Human Molecular Genetics, 4, pp
1821-1827
11. Gius et al., (1990), Mol. Cell. Biol., 10, pp 4243-4255
12. Gurney et al., (1992), J. Biol. Chem., 267, pp 18133-
18139
- 20 13. Hastay et al., (1990), Arthritis Rheum., 33, pp 388-397
14. Hay et al., (1989), Genes Dev., 3, pp 293-303
15. Heuertz et al., (1993), Genomics, 18, pp 100-104
16. Holt et al., (1986), Proc. Natl. Acad. Sci. USA, 83,
pp 4794-4798
- 25 17. Hu et al., (1994), EMBO J., 13, pp 3094-3103
18. Keck et al., (1989), Science, 246, pp 1309-1311
19. Kerr et al., (1988), Science, 242, pp 1424-1427
20. Kim et al., (1993), Nature, 362, pp 841-844
21. Kovary & Bravo, (1991), Mol. Cell. Biol., 11, pp 2451-
30 2459
22. Kovary & Bravo, (1991), Mol. Cell. Biol., 11, pp 4466-
4472
23. Leung et al., (1989), Science, 246, pp 1306-1309
24. Liang et al., (1993), NAR, 21, pp 3269-3275
- 35 25. Liotta and Stetler, (1990), Semin. Cancer Biol., 1, pp
99-106
26. Lord et al., (1993), Mol. Cell. Biol., 13, pp 841-851
27. Miller et al., (1984), Cell, 36, pp 51-60

28. Plate et al., (1993) Cancer Research, 53, pp 5822-5827
29. Riabowol et al., (1988), Mol. Cell. Biol., 8, pp 1670-1676
30. Rollins et al., (1988), Proc. Natl. Acad. Sci. USA, 85, pp 3738-3742
- 5 31. Ruther et al., (1989), Oncogene, 4, pp 861-865
32. Sambrook et al., (1989), Molecular Cloning: A Laboratory Manual. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- 10 33. Sassone et al., (1988), Nature, 334, pp 314-319
34. Schonthal et al., (1988), Cell, 54, pp 325-334
35. Schusled et al., (1995), Brain Res., 670, pp 14-28
36. Seed & Aruffo, (1987), Proc. Natl. Acad. Sci. USA, 84, pp 3365-3369
- 15 37. Superti-Furga et al., (1991), Proc. Natl. Acad. Sci. USA, 88, pp 5114-5118
38. Takeshita et al., (1994), J. Clin. Invest., 93, pp 662-670
39. Tsunoda et al., (1994), Anti-cancer Res., 14, pp 2637-2642
- 20 40. Vaziri et al., (1995), Mol. Cell. Biol., 15, pp 1244-1253
41. Woessner and Gurja,, (1991), J. Rheumatol., Suppl. 27, pp 99-101
- 25 42. EP-A-0120693
43. EP-A-0125023

Claims

1. A nucleotide molecule encoding a protein encoded by a
5 Fos regulated gene or a fragment thereof, wherein said
protein or fragment thereof is encoded by any one of the
nucleotide sequences shown in Figure 1 or 2 or a fragment
thereof, including allelic variants and species variants of
the nucleotide sequences.
- 10 2. The nucleotide molecule of claim 1, wherein said protein
or a fragment thereof, encoded by the sequences shown in
Figure 1 or 2 is altered but still has at least 80% homology
to the protein or a fragment thereof, encoded by the
15 sequences shown in Figure 1 or 2.
3. The protein encoded by the nucleotide molecule of claim
1 or claim 2.
- 20 4. A vector for the expression of the nucleotide molecule
of claim 1 or claim 2, comprising a promoter and said
nucleotide molecule.
- 25 5. A host cell transformed with the vector of claim 4.
6. The host cell of claim 5 which is a chinese hamster
ovary cell.
7. A method for the producing the protein of claim 3,
30 comprising culturing the host cell of claim 5 or claim 6
under conditions leading to the production of the protein
and harvesting the protein.
8. The nucleotide molecule of claim 1 or claim 2 for use
35 in therapy.
9. The use of the nucleotide molecule of claim 1 or claim
2 in the manufacture of a composition for the treatment of

developmental disorders.

10. An antibody molecule having specificity for the protein of claim 3.

5

11. The antibody molecule of claim 10 for use in therapy.

12. The use of the antibody molecule of claim 10 in the manufacture of a composition for the treatment of
10 proliferative diseases.

13. An antisense nucleotide molecule having the complementary sequence to the nucleotide molecule of claim 1 or claim 2.

15

14. An antisense vector for the expression of the antisense nucleotide molecule of claim 13, comprising a promoter and the antisense molecule.

20 15. The antisense vector of claim 14 for use in therapy.

16. The use of the antisense vector of claim 14 in the manufacture of a composition for the treatment of proliferative diseases.

25

17. A vector for the expression of a ribozyme, comprising a promoter and a nucleotide sequence encoding a ribozyme capable of cleaving the RNA transcript of the nucleotide molecule of claim 1 or claim 2.

30

18. The vector of claim 17 for use in therapy.

19. The use of the vector of claim 17 in the manufacture of a composition for the treatment of proliferative diseases.

35

20. The use of claim 12, 16 or 19, wherein the proliferative disease is cancer.

21. The use of the protein of claim 3 in identifying the receptor of said protein.
22. The use of the protein of claim 3 in an assay for the
5 identification of antagonists or agonists of said protein.
23. The use of the nucleotide molecule of claim or claim 2, the protein of claim 3 or the antibody molecule of claim 10
10 in diagnosing a pathological state or a predisposition to a disease.
24. The use of the nucleotide sequence of claim 1 or claim 2 in the generation of a transgenic animal.
- 15 25. A nucleotide molecule comprising all or part of the sequence shown in any one of Figures 1 or 2.

1/31

FIG. 1

```

10      30      50
ggaagatatgaccacctcctgattattttgacgagggaaccttgaaatatattttcatt
.      .      .
70      90      110
gctttctccatactaagattgtgtgaggcagtgaggagtccttgacttactcaag
.      .      .
130     150     170
tcatttcattggattttaattacaactgatcatgtgattgttttttccatgtaaagttt
.      .      .
190     210     230
ggggcttcaaaactttgcttctggagaatgccttttgcaacaccttttcagtagctgcctgg
.      .      .
250     270     290
aaacaactgcttagtcatcggtagacattttaaataattcaaaatgtatggagaatgggga
.      .      .

```

M Y G E W G

2/31

FIG. 1(1)

310	330	350
atggggaatacctcatgatgttccatgtgtacttgggtgcagggttcaggagcgaacat		
M G N I L M M F H V Y L V Q G F R S E H		
370	390	410
ggaccagtgaaggatttttctttttgagcgcgcacatcccgggtccatgttggaacgatctgaa		
G P V K D F S F B R S S R S M L E R S E		
430	450	470
caacagatccgagcagcttctagtttgaggagtgctgcaaatcgcgcactctgaggac		
Q Q I R A A S S L E B L L Q I A H S E D		

3/31

FIG. 1(III)

490	510	530
tggaagctgtggcgatgccggttgaagctcaaaagtcttgccagtatggactcacgctca		
W K L W R C R L K L K S L A S M D S R S		
550	570	590
gcattcccatcgctccaccagatttgcggcaactttcttatgacactgaaacactaaaagtt		
A S H R S T R F A A T F Y D T E T L K V		
610	630	650
atagatgaagaatggcagaggaccaatgacagccctagagagacatgcgtagaagtcgcc		
I D E E W Q R T Q C S P R E T C V E V A		
670	690	710
agtgagctggggaagacaaccaacacattottcaagccccctgtgtaaatgtcttcgg		
S E L G K T T N T F F K P P C V N V F R		

FIG. 1 (iii)

730	750	770
tgtggaggctgctgcaacgaagagggtgtgtgtgtatgtaacaagaacctctacatc		
cgcgcgcncnecgvmcmntstsyi		

790 810 830
tccaaacagctctttgagatatcagtgccctctgacatcagtgcccgagttagtgccctgtt
S K Q L F E I S V P L T S V P E L V P V

850
870
890
aaaattgccaaaccatacgggttgtaagtgttgccacggggcccccattctctca
K I A N H T G C K C L P T G P R H P Y S

attatcagaagatccattcagacccagagaagatgaatgtctcattccaagaaactc
I I R R S I Q T P E R D E C P H S K K L

	970		990	1010															
C	P	I	D	M	L	W	D	N	T	K	C	K	C	V	L	Q	D	E	T
tgtcctattgacatgc	tgtgggataaaccca	aattgtaaatgtgttttgc	aagacgagact																

FIG. 1(IV)

1030	1050	1070
ccactgcctgggacagaagaccactcttaccctccaggaaccactctctgtggaaccgcac P L P G T E D H S Y L Q E P T L C G P H		
1090	1110	1130
atgacgcttgatgaagatcgctgtgagtgcgctctgtaaagcaccatgtccgggagatctc M T F D E D R C E C V C K A P C P G D L		
1150	1170	1190
attcagcaccggaaaactgcagttgcttttgagtgcaagaagaaagtctggagagctgctgc I Q H P E N C S C F E C K E S L E S C C		
1210	1230	1250
caaaagcacaagattttcaccacagacacctgcagctgtgaggacagatgtccttttcac Q K H K I F H P D T C S C E D R C P F H		
1270	1290	1310
accagaacatgtgcaagtagaaagccagcctgtggaagcactggcgctttccaaggag T R T C A S R K P A C G K H W R F P K E		

5/31

FIG. 1(V)

1330 1350 1370
 . . .
 acaagggccaggactctacagccaggagaaacccttgattcaacttccttcaagtccc
 T R A Q G L Y S Q E N P

 1390 1410 1430
 . . .
 cccatctctgtcattttaaacagctcactgctttgtcaagttgctgtcactgttgccac

 1450 1470 1490
 . . .
 taccctgccccccccctccccgcctccagggtgttagaaaagttgatttgacctagtgt
 6/31

 1510 1530 1550
 . . .
 catggtaagccacattttccatgcaatggcggctaggtagtattccccagttcactgacaaa

 1570 1590 1610
 . . .
 tgacttgtagcttcagatgtctttgcgccatcagcactcagaaaaggaaggggtctgagga

7/31

1630	1650	1670
gccccttggtttgatgaataagaaagggttgccctgaaacagagtagtaggtgccactcga		
1690	1710	1730
ttgggttcctcgggctggcaagtcgaagggaatgctcatgagttattgtgcttcttctt		
1750	1770	1790
tatgcggaatttcatttgatgatcagcactgatcaattccattccacttgacttttt		
1810	1830	1850
aggttactgaagcactgcctgatgttttatatgtaaatgtatttaagggaataaacac		
1870	1890	
tgttatgcagcccaaaaaaaaaa		

FIG. 1(VI)

FIG. 2

8/31

```

1  GGCACGAGGTTTTTTTTTTTTTTCATCTCTCTCTCCCAACCCTAAGATTGTGCAA
   +-----+-----+-----+-----+-----+-----+
60 CCGTGCTCCAAAAAAGTAGAGAGAGAGGGGTGGGATTTCTAACACGTTT
   +-----+-----+-----+-----+-----+-----+
   G T R F F F F F I S L S P P L R L C K -
   A R G F F F F S S L S P H P * D C A K -
   H E V F F F F H L S L P T P K I V Q K -
   +-----+-----+-----+-----+-----+-----+
61 AAAAGCGTACCTTGCCTAATTGAATAATTTCATTGGATTTCAGAACTGATTATT
   +-----+-----+-----+-----+-----+-----+
120 TTTTCGCATGGAACGGATTAACTTTATTAAGTAACCTAAACTAGTCTTGACTAATAAA
   +-----+-----+-----+-----+-----+-----+
   K S V P C L I E I I S L D F D Q N * L F -
   K A Y L A * L K * F H W I L I R T D Y L -
   K R T L P N * N N F I G F * S E L I I W -
   +-----+-----+-----+-----+-----+-----+
121 GGTTTCTGTGAAGTTTTCAGGTTTCAAACTTTCCTTCTGGAGAAATGCCTTTGTGAAC
   +-----+-----+-----+-----+-----+-----+
180 CCAAAGACACACTTCAAAACCTCCAAAGTTTGAAGGAAGACCTCTTACGGAAACTTTG
   +-----+-----+-----+-----+-----+-----+

```

FIG. 2(1)

G F L C E V L R F Q T F L L E N A F * N -
 V F C V K F * G F K L S F W R M P F E T -
 F S V * S F E V S N F P S G E C L L K Q -
 AATTTCCTAGCTGCCCTGATGTCACCTGCTTAGTAATCAGTGGATATTGAAATATTCAA + 240
 TTAAAGAGATCGACGACTACAGTTGACCAATCATTAGTCACCTATTAACCTTATAAGTT
 N F L * L P D V N C L V I S G Y * N I Q -
 I F S S C L M S T A * S V D E E I F K -
 F S L A A * C Q L L S N Q W I L K Y S K -
 AATGACAGAGTGGTAGTGGTGAATGTTTCATGATGTTGTACGTCACGCTGTGCA + 300
 TTACATGCTCTCACCCTACCACTTACAAAGTACTACAACATGCAGTCGACCCACGT
 N V Q R V G S G E C E H D V V R P A G A -
 M Y R E W V V N V F M M L V V Q L V Q -
 C T E S G * W * M F S * C C T S S W C R -

9/31

FIG. 2(II)

301 GGGCTCCAGTAATGAACATGGACCAGTGAAGCCGATCATCTCAGTCCACATTGGAACGATC
+-----+-----+-----+-----+-----+-----+
360 CCCGAGGTCATTACTTGTACTGCTGGTCACTTCGCTAGTAGAGTCAGGTGTAACCTTGCTAG

G L Q * * T W T S E A I I S V H I G T I -
[G S S N E H G P V K R S S Q S T L E R S] -
A P V M N M D Q * S D H L S P H W N D L -

361 TGAACAGCAGATCAGGGCTGCTTCTAGTTGGAGGAACTACTTCGAAATTACTCACTCTGA
+-----+-----+-----+-----+-----+-----+
420 ACTTGTCTAGTCCCGACGAGATCAAACTCCTTGATGAAGCTTAATGAGTGAGACT

* T A D Q G C F * F G G T T S N Y S L * -
[E Q Q I R A A S S L E E L L R T T H S E] -
N S R S G L L L V W R N Y F E L L T L R -

GGACTGGAAGCTGTGAGATGCAGGCTGAGGCTCAAAGTTTACCAGTATGGACTCTCG

FIG. 2(III)

421 -----+-----+-----+-----+-----+-----+-----+ 480
 CCTGACCTTCGACACCTCTACGTCCGACTCCGAGTTTCAAAATGGTCATACCTGAGAGC
 L E A V E M Q A E A Q K F Y Q Y G L S -
 [D W K L W R C R L R L K S F T S M D S R]
 T G S C G D A G * G S K V L P V W T L A -
 CTCAGCATCCCATCGGTCCACTAGGTTGGGGCAACTTCTATGACATTGAAACACTAAA 540
 -----+-----+-----+-----+-----+-----+-----+
 GAGTCGTAGGGTAGCCAGGTGATCCAAACGCCGTTGAAGATACCTGTAACTTGTGATTT
 L S I P S V H * V C G N F L * H * N T K -
 [S A S H R S T R F A A T F V D I E T L K]
 Q H P I G P L G L R Q L S M T L K H * K -
 AGTTATAGATGAAGAATGGCAAGAACTCAGTGCAGCCCTAGAGAAACGTGGAGGT 600
 -----+-----+-----+-----+-----+-----+-----+
 TCAATATCTACTTCTTACCCTTCTTNGAGTCACGTCCGGATCTCTTTGCACGCACCTCCA
 S Y R * R M A K N S V Q P * R N V R G G -
 [V I D E E W Q R T Q C S P R E T C V E V]
 L * M K N G K E L S A A L E K R A W R W -

///31

FIG. 2(IV)

601 GGCAGTGAGCTGGGAAGAGTACCAACACATTCTTCAAGCCCCCTTGTGTGAACGTGTT
 660 CCGTCACTCGACCCCTTCTCATGGTTGTGTGAAGAAGTTCCGGGGAACACACTTGCACAA

G Q * A G E E Y Q H I L Q A P L C E R V
 A S E L G K S T N T F F K P P C V N V E
 P V S W G R V P T H S S S P L V * T C S

661 CCGATGTGGTGGTGTGCAATGAAGAGAGCCTTATGTGTATGAACACCAGCACCTCGTA
 720 GGCTACACCACCAACGTTACTTCTCTCGAATAACACATACATTGTGTCGTGGAGCAT

P M W W L L Q * R E L Y V Y E H Q H L V
 R C G G C C N E E S F M C M N T S T S Y
 D V V V V A M K R A L C V * T P A P R T

721 CATTCCAAACAGCTCTTTGAGATATCAGTGCCCTTGCATCAGTACCTGAATTAGTCC
 780 GTAAAGGTTTGTGAGAAACTCTATAGTCACGGAAACTGTAGTCATGGACTTAATCACGG

H F Q T A L * D I S A F D I S T * I S A
 I S K Q L F E I S V P L T S V P E L V P
 F P N S S L R Y Q C L * H Q Y L N * C L

FIG. 2(v)

781 TGTAAAGTTGCCAATCATACAGGTTGTAAGTCTTGCCAAACAGCCCCCGCCATCCATA
 840 ACAATTCAACGGTTAGTATGTCCTCAACATTACGAAACGGTTGTCGGGGGGCGGTAGGTAT

C * S C Q S Y R L * V L A N S P P S I -
 V K V A N H T G C K C L P T A P R H P Y -
 L K L P I I Q V V S A C Q Q P P A I H T -

841 CTCATTATCAGAAGATCCATCCAGATCCCTGAAGAAGATCGCTGTCCCATCCAAAGAA
 900 GAGTTAATAGTCTTCTAGGTAGGTCTAGGACTTCTTCTAGCCGACAAGGTAAGGTTCTT

L N Y Q K I H P D P * R R S L F P F Q E -
 S I I R R S I Q I P E E D R C S H S K K -
 Q L S E D P S R S L K K I A V P I P R N -

ACTCTCTCTATTGCATGCTATGGATACCAACAAATGTAATGTGTTTGCAGGAGGA

13/31

15/31

FIG. 2(vii)

1081	TCATAATCCAGCACCCCAAAACTGCAGTTGCTTTGAGTGCAAGAAGTCTGGAGACCTG	-
	AGATTAGGTCGTGGGCTTTTGACGTCAACGAACCTCACGTTTCTTTCAGACCTCTGGAC	1140
	S N P A P Q K L Q L L * V Q R K S G D L	-
	L I Q H P K N C S C F E C K E S E E T C	-
	* S S T P K T A V A L S A K K V W R P A	-
1141	CTGCCAGAAGCACAAGCTATTTCACCCAGACACCTGCAGCTGTGAGGACAGATGCCCTT	-
	GACGGCTTCGTGTTGATAAAGTGGTCTGTGGACGTCCACACTCCTGTCTACGGGGA	1200
	L P E A Q A I S P R H L Q L * G Q M P L	-
	C Q K H K L F H P D T C S C E D R C P F	-
	A R S T S Y F T Q T P A A V R T D A P F	-
1201	TCATACCAGACCATGTGCAAGTGGCAAAACAGCATGTGCAAGCATGCCGCTTCCAA	-
	AGTATGGTCTGGTACACGTTACCGTTTGTGCTACACGTTTCGTAAACGGCGAAAGGTTT	1260
	S Y Q T M C K W Q N S M C K A L P L S K	-
	H T R P C A S G K T A C A K H C R F P K	-
	I P D H V Q V A K Q Q H V Q S I A A F Q R	-

FIG. 2(viii)

1261 GGAGAAAGGGCTGCCAGGGCCCCACAGCCGAAAGAAATCCTTGA^{16/31}TTCAGCGTTCCAAG 1320
 -----+-----+-----+-----+-----+-----+-----+
 CCTCTTTCCCGACGGGTCCCCGGGTGTCGGCTTCTTAGGAACATAAGTCGCAAGGTTTC
 G E K G C P G A P Q P K E S L I Q R S K -
 E K R A A Q G P H S R K N P * F S V P S -
 R K G L P R G P T A E R I L D S A F Q V -
 1321 TTCCCATCCCTGTCA^{16/31}TTTAAACAGCATGCTGCTTTGCCAAGTTGCTGTCACTGTTT 1380
 -----+-----+-----+-----+-----+-----+-----+
 AAGGGTAGGGACAGTAA^{16/31}AAATGTCTACGACGAAACGGTTCAACGACAGTGACAA^{16/31}AA
 F P I P V I F N S M L L C Q V A V T V F -
 S P S L S F L T A C C F A K L L S L F F -
 P H P C H F * Q H A A L P S C C H C F F -
 TTCCCACTGTAA^{16/31}AAAAAATCCATTTTACACAGCACCACAGTGAATCCACACCAACC

FIG. 2(ix)

1381 -----+-----+-----+-----+-----+-----+-----+-----+-----+-----+ 1440
 AAGGTCACAATTTTTTTAGGTAAATGTCGTGGTGCACCTTAGGTCGTGGTTGG
 F P G V K K K I H F T Q H H S E S R P T -
 S Q V L K K K S I L H S T T V N P D Q P -
 P R C * K K N P F Y T A P Q * I Q T N L -
 1441 -----+-----+-----+-----+-----+-----+-----+-----+-----+-----+ 1500
 TTCCATTACACCAGCTAAGGAGTCCCTGGTTTCATTGATGGATGCTTCTTAGCTGCAGAT
 AAGGTAAGTGTGGTCGATTCTCTCAGGACCAAGTAACCTACAGAAAGATCGACGTCTA
 F H S H Q L R S P W F I D G C L L A A D -
 S I H T S * G V P G S L M D V F * L Q M -
 P F T P A K E S L V H * W M S S S C R C -
 1501 -----+-----+-----+-----+-----+-----+-----+-----+-----+-----+ 1560
 GCCTCTGCGCACCAAGGAATGGAGAGGGGACCCATGTAATCCTTTTGTAGTTTIG
 CGGAGACGCGTGGTTCCTTACCTCTCCTCCCTGGGTACATTAGGAAACAAATCAAAC
 A S A H Q G M E R R G P M * S F C L V L -
 P L R T K E W R G G D P C N P F V * F C -
 L C A P R N G E E G T H V I L L F S F V -

17/31

FIG. 2(x)

1561 TTTTGTGTTTGGTGAATGAGAAAGGTGTGCTGTCATGGAATGGCAGGTGCATATGA
 -----+-----+-----+-----+-----+
 1620 AAAACAAAAACCCTTACTCTTTCCACAGACCAGTACCTTACCGTCCACAGTATACT
 -----+-----+-----+-----+-----+
 F L F F G E * E R C A G H G M A G V I *
 F C F L V N E K G V L V M E W Q V S Y D
 F V F W * M R K V C W S W N G R C H M T
 -----+-----+-----+-----+-----+
 1621 CTGATTACTCAGAGCAGATGAGGAAACTGTAGTCTCTGAGTCCTTTGCTAATCGCAACT
 -----+-----+-----+-----+-----+
 1680 GACTAATGAGTCTGTCCTTCTACTCTTTTGACATCAGAGACTCAGGAACGATTAGCGTTGA
 -----+-----+-----+-----+-----+
 L I T Q S R * G K L * S L S P L L I A T
 * L L R A D E E N C S L * V L C * S Q L
 D Y S E Q M R K T V V S E S F A N R N S
 -----+-----+-----+-----+-----+
 1681 CTGIGAATTATTCTGATTCTTTTATGCAGAAATTGATTGCTATGATCAGTACTGACT
 -----+-----+-----+-----+-----+
 1740 GAACACTTAATAAGACTAAGGAAATAACGTCTTAACTAAGCATACTAGTCATGACTGA
 -----+-----+-----+-----+-----+
 L V N Y S D S F L C R I * F V * S V L T
 L * I I L I L F Y A E F D S Y D Q Y * L
 C E L F * F F F M Q N L I R M I S T D F
 -----+-----+-----+-----+-----+

18/31

1741 TTCTGATTACIGTCCAGCCTTATAGTCTTCCAGTTTAAATGAACCTACCATCTGATGTTTCAT
 -----+-----+-----+-----+-----+-----+-----+
 AAGACTAATGACAGGTGCGAATATCAGAAGGTCAAAATTACTTGATGGTAGACTACAAAGTA
 -----+-----+-----+-----+-----+-----+-----+ 1800
 F * L L S S L * S S S L M N Y H L M F H -
 S D Y C P A Y S L P V * * T T I * C F I -
 L I T V Q L I V F Q F N E L P S D V S Y -
 1801 ATTTAAGTGTATTAAAGAAATAAACACCATTTATTCAGCCATATAAATAAATAAATAA
 -----+-----+-----+-----+-----+-----+-----+ 1860
 TAAATTCACATAAATTCTTTTATTGTGGTAATAAGTTCGGTATATTTTITTTTTTTT
 I * V Y L K K I N T I I Q A I * K K K K -
 F K C I * R K * T P L F K P Y K K K K -
 L S V F K E N K H H Y S S H I K K K K -
 AAAA

FIG. 2(xi)


20/31

FIG. 3


FDGF	(103)	TLKVIDEEWQ	•	RTQCSPRETC	VEVASELGKT	T..NTFFKPP	•	CVNVFRCGGC
VEGF-C	(49)	ILKSIDNEWR		KTCMPREVC	IDVGKEFGVA	T..NTFFKPP		CVSVYRCGGC
VEGF	(39)	EVVKFMDVYQ		RSYCHPIETL	VDIFQEYPDE	I..EYIFKPS		CVPLMRCGGC
PIGF	(39)	EVVPFQEVWG		RSYCRALERL	VDVSEYPSE	V..EHMFSPS		CVSLLRCTGC
PDGF-B	(84)	GSLTIAEPAM		IAECKTRTEV	FEISRRRLDR	TNANFLVWPP		CVEVQRCSCG
PDGF-A	(83)	RRKRSIEEAV		PAVCKTRTVI	YEIPRSQVDP	TSANFLIWPP		CVEVKRCTGC
FDGF	(151)	• CNEEGVMCMN		TSTSISKQL	FEIS.VPLTS	VPELVPVKIA	•	NHTGCKCLPT
VEGF-C	(97)	CNSEGLQCMN		TSTSLSKTL	FEIT.VPLSQ	GPKPVTISFA		NHTSCRCMSK
VEGF	(87)	CNDEGLECVP		TEESNITMQI	MRIK..P.HQ	GQHIGEMSFL		QHNKCECRPK
PIGF	(87)	CGDENLHCVP		VETANVTMQL	LKIR..S.GD	RPSYVELTFS		QHVRCECRPL
PDGF-B	(134)	CNNRNVCQCP		TQVQLRPVQV	RKIEIVRKKP	IFKKATVTLE		DHLACKCETV
PDGF-A	(133)	CNTSSVKCQCP		SRVHRSVKV	AKVEYVRKKP	KLKEVQVRLE		EHLECACATT

21/31


FIG. 4(A)



22/31


**FIG. 4(B)**

23/31




SUBSTITUTE SHEET (RULE 26)

24/31


**FIG. 4(D)**

25/31

**FIG. 4(E)**


26/31

FIG. 5(A)




27/31

FIG. 5(B)



28/31

FIG. 6



29/31

FIG. 7(C)





FIG. 7(B)



FIG. 7(A)



30/31

FIG. 7(F)




FIG. 7(E)





FIG. 7(D)



31/31

FIG. 7(G)





FIG. 7(H)



SUBSTITUTE SHEET (RULE 26)

THIS PAGE BLANK (USPTO)



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/19, 5/10, C07K 14/52, 16/24, C12Q 1/68, G01N 33/574, A61K 31/70		A3	(11) International Publication Number: WO 97/12972
			(43) International Publication Date: 10 April 1997 (10.04.97)
(21) International Application Number: PCT/IB96/01113		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 30 September 1996 (30.09.96)		Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(30) Priority Data: 9519928.7 29 September 1995 (29.09.95) GB 9612368.2 13 June 1996 (13.06.96) GB			
(71) Applicant (for all designated States except US): UNIVER- SITA'DEGLI STUDI DI SIENA [IT/IT]; Via Banchi di Sotto, 55, I-53100 Siena (IT).			
(72) Inventor; and (75) Inventor/Applicant (for US only): OLIVIERO, Salvatore [IT/IT]; Via del Giglio, 20, I-53100 Siena (IT).			
(74) Agent: HALLYBONE, Huw, George; Carpmacels & Ransford, 43 Bloomsbury Square, London WC1A 2RA (GB).		(88) Date of publication of the international search report: 12 June 1997 (12.06.97)	
(54) Title: REGULATED GENES AND USES THEREOF			
(57) Abstract			
<p>A nucleotide molecule encoding a protein encoded by a Fos regulated gene or a fragment thereof, wherein said protein or fragment thereof is encoded by any one of the nucleotide sequences shown in Figure 1 or 2 or a fragment thereof, including allelic variants and species variants of the nucleotide sequences.</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LJ	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

INTERNATIONAL SEARCH REPORT

International Application No

PCT/IB 96/01113

A. CLASSIFICATION OF SUBJECT MATTER
 IPC 6 C12N15/19 C12N5/10
 G01N33/574 A61K31/70

C07K14/52

C07K16/24

C12Q1/68

According to International Patent Classification (IPC) or to both national classification and IPC.

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N C07K C12Q G01N A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EMBL SEQUENCE DATABASE, 9 July 1995, HEIDELBERG, BRD, XP002029337 L. HILLIER ET AL.: "y142g02.ri Homo sapiens cDNA clone 160946 5'" see the whole document ---	25
A	EMBO J., vol. 13, no. 13, 1 July 1994, OXFORD UNIVERSITY PRESS, GB;; pages 3094-3103, XP002029338 E. HU ET AL.: "Targeted disruption of the c-fos gene demonstrates c-fos-dependent and -independent pathways for gene expression stimulated by growth factors or oncogenes" cited in the application see the whole document --- -/-	1-25



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

G document member of the same patent family

Date of the actual completion of the international search

14 April 1997

Date of mailing of the international search report

18.04.97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
 Fax (+ 31-70) 340-3016

Authorized officer

Hornig, H

INTERNATIONAL SEARCH REPORT

International Application No

PCT/IB 96/01113

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>NATURE, vol. 362, 29 April 1993, MACMILLAN JOURNALS LTD., LONDON, UK, pages 841-844, XP002029339 K. KIM ET AL.: "Inhibition of vascular endothelial growth factor-induced angiogenesis supresses tumour growth in vivo" cited in the application see the whole document ---</p>	1-25
A	<p>SCIENCE, vol. 246, no. 4935, 8 December 1989, pages 1306-1309, XP000083690 LEUNG D W ET AL: "VASCULAR ENDOTHELIAL GROWTH FACTOR IS A SECRETED ANGIOGENIC MITOGEN" cited in the application see the whole document ---</p>	1-25
A	<p>SCIENCE, vol. 246, no. 4935, 8 December 1989, pages 1309-1312, XP000083691 KECK P J ET AL: "VASCULAR PERMEABILITY FACTOR, AN ENDOTHELIAL CELL MITOGEN RELATED TO PDGF" cited in the application see the whole document ---</p>	1-25
T	<p>PROC. NATL.ACAD SCI., vol. 93, no. 21, 15 October 1996, NATL. ACAD SCI., WASHINGTON, DC, US., pages 11675-11680, XP002029340 M. ORLANDINI ET AL.: "Identification of a c-fos induced gene that is related to the platelet-derived growth factor/ vascular endothelial growth factor family" see the whole document -----</p>	1-25

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ **BLACK BORDERS**

☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**

☐ **FADED TEXT OR DRAWING**

☒ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**

☐ **SKEWED/SLANTED IMAGES**

☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**

☐ **GRAY SCALE DOCUMENTS**

☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**

☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**

☐ **OTHER: _____**

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.